

University of Groningen

## The adaptive value of migrations for the bivalve *Macoma balthica*

Hiddink, Johan Gerrit

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2002

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Hiddink, J. G. (2002). *The adaptive value of migrations for the bivalve Macoma balthica*. s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# REFERENCES

---

- Aarnio K, Bonsdorff E, & Norkko A (1998) Role of *Halicryptus spinulosus* (Priapulida) in structuring meiofauna and settling macrofauna. *Marine Ecology Progress Series*, 163, 145-153.
- Abrahams M & Kattenfeld M (1997) The role of turbidity as a constraint on predator-prey interactions in aquatic environments. *Behavioral Ecology and Sociobiology*, 40, 169-174.
- Adema JPHM (1991) *De krabben van Nederland en België*. Nationaal Natuurhistorisch Museum, Leiden.
- Afman BP (1980) De voedelopname van de strandkrab *Carcinus maenas* (L.). *Interne Verslagen Nederlands Instituut voor Onderzoek der Zee*, 1980-4, 1-28.
- Ambrose WG (1984a) Influences of predatory polychaetes and epibenthic predators on the structure of a soft-bottom community in a Maine estuary. *Journal of Experimental Marine Biology and Ecology*, 130, 167-187.
- Ambrose WG (1984b) Role of predatory infauna in structuring marine soft-bottom communities. *Marine Ecology Progress Series*, 17, 109-115.
- Anderson A (1971) Intertidal activity, breeding and the floating habit of *Hydrobia ulvae* in the Ythan estuary. *Journal of the Marine Biological Association of the United Kingdom*, 51, 423-437.
- Andre C & Rosenberg R (1991) Adult-larval interactions in the suspension-feeding bivalves *Cerastoderma edule* and *Mya arenaria*. *Marine Ecology Progress Series*, 71, 227-234.
- Ansell AD, Barnett PRO, & Massé H (1981) Upper temperature tolerances of some European molluscs. III. *Cardium glaucum*, *C. tuberculatum* and *C. edule*. *Marine Biology*, 65, 177-183.
- Ansell AD, Comely CA, & Robb L (1999) Distribution, movements and diet of macrocrustaceans on a Scottish sandy beach with particular reference to predation on juvenile fishes. *Marine Ecology Progress Series*, 176, 115-130.
- Armonies W (1992) Migratory rhythms of drifting juvenile molluscs in tidal waters of the Wadden Sea. *Marine Ecology Progress Series*, 83, 197-206.
- Armonies W (1994a) Drifting meio- and macrobenthic invertebrates on tidal flats in Königshafen: a review. *Helgoländer Meeresuntersuchungen*, 48, 299-320.
- Armonies W (1994b) Turnover of postlarval bivalves in sediments of tidal flats in Königshafen (German Wadden Sea). *Helgoländer Meeresuntersuchungen*, 48, 291-297.
- Armonies W (1996) Changes in distribution patterns of 0-group bivalves in the Wadden Sea: Byssus-drifting releases juveniles from the constraints of hydrography. *Journal of Sea Research*, 35, 323-334.
- Armonies W (2001) What an introduced species can tell us about the spatial extension of benthic populations. *Marine Ecology Progress Series*, 209, 289-294.
-

- Armonies W & Hartke D (1995) Floating of mud snails *Hydrobia ulvae* in tidal waters of the Wadden Sea, and its implications in distribution patterns. *Helgoländer Meeresuntersuchungen*, 49, 529-538.
- Armonies W & Hellwig-Armonies M (1992) Passive settlement of *Macoma balthica* spat on tidal flats of the Wadden Sea and subsequent migration of juveniles. *Netherlands Journal of Sea Research*, 29, 371-378.
- Bachelet G (1980) Growth and recruitment of the tellinid bivalve *Macoma balthica* at the southern limit of its geographical distribution, the Gironde estuary (SW France). *Marine Biology*, 59, 105-117.
- Baker RR (1978) *The evolutionary ecology of animal migration*. Hodder and Stoughton, London.
- Barber VC & Land MF (1967) Eye of the cockle, *Cardium edule*: anatomical and physiological observations. *Experientia*, 23, 677-678.
- Barnes RSK (1981) Behavioural activities and ecological strategies in the intertidal gastropod *Hydrobia ulvae*. *Feeding and survival strategies of estuarine organisms*. (eds. N. V. Jones & W. J. Wolff), pp. 79-90 Plenum press, New York and London.
- Barnes RSK (1998) The effects of movement on population density estimates of mudflat epifauna. *Journal of the Marine Biological Association of the United Kingdom*, 78, 377-385.
- Bayne BL (1964) Primary and secondary settlement in *Mytilus edulis* L. (Mollusca). *Journal of Animal Ecology*, 33, 513-523.
- Beaumont AR & Barnes DA (1992) Aspects of veliger larval growth and byssus drifting of the spat of *Pecten maximus* and *Aequipecten opercularis*. *ICES Journal of Marine Science*, 49, 417-423.
- Benfield MC & Minello TJ (1996) Relative effects of turbidity and light intensity on reactive distance and feeding of an estuarine fish. *Environmental Biology of Fishes*, 46, 211-216.
- Benham WB (1893) The early post-larval stage of *Arenicola marina*. *Journal of the Marine Biological Association of the United Kingdom*, 3, 47-52.
- Berge JA & Hesthagen IH (1981) Effects of epibenthic macropredators on community structure in an eutrophicated shallow water area, with special reference to food consumption by the common goby *Pomatoschistus microps*. *Kieler Meeresforschung*, 5, 462-470.
- Berger VY & Khalaman VV (1996) The non-spawning migration of polychaete *Nereis virens* Sars at the White Sea. *Doklady Akademii Nauk*, 349, 710-711.
- Berry AJ (1988) Annual cycle in *Retusa obtusa* (Montagu) (Gastropoda, Ophisthobranchia) of reproduction, growth and predation upon *Hydrobia ulvae* (Pennant). *Journal of Experimental Marine Biology and Ecology*, 117, 197-209.
- Berthold P (1993) The phenomena of bird migration. *Bird migration, a general survey*. (ed. P. Berthold), pp. 31-86 Oxford university press, Oxford.
- Beukema JJ (1985) Zoobenthos survival during severe winters on high and low tidal flats in the Dutch Wadden Sea. *Marine biology of polar regions and effects of stress on marine organisms*. (eds. J. S. Gray & M. E. Christiansen), pp. 351-361 John Wiley, Chichester.
- Beukema JJ (1991) The abundance of shore crabs *Carcinus maenas* (L.) on a tidal flat in the Wadden Sea after cold and mild winters. *Journal of Experimental Marine Biology and Ecology*, 153, 97-114.
- Beukema JJ (1992) Dynamics of juvenile shrimp *Crangon crangon* in a tidal-flat nursery of the Wadden Sea after mild and cold winters. *Marine Ecology Progress Series*, 83, 157-165.
- Beukema JJ (1993a) Successive changes in distribution patterns as an adaptive strategy in the bivalve *Macoma balthica* (L.) in the Wadden Sea. *Helgoländer Meeresuntersuchungen*, 47, 287-304.
- Beukema JJ (1993b) Increased mortality in alternative bivalve prey during a period when the tidal flats of the Dutch Wadden Sea were devoid of mussels. *Netherlands Journal of Sea Research*, 31, 395-406.
- Beukema JJ (1997) Caloric values of marine invertebrates with an emphasis on the soft parts of marine bivalves. *Oceanography and Marine Biology: an annual review*, 35, 387-414.
- Beukema JJ & De Vlas J (1979) Population parameters of the lugworm, *Arenicola marina*, living on tidal flats in the Dutch Wadden Sea. *Netherlands Journal of Sea Research*, 13, 331-353.

- 
- Beukema JJ & De Vlas J (1989) Tidal-current transport of thread-drifting postlarval juveniles of the bivalve *Macoma balthica* from the Wadden Sea (Netherlands) to the North Sea. *Marine Ecology Progress Series*, 52, 193-200.
- Beukema JJ, Essink K, & Michaelis H (1996) The geographic scale of synchronized fluctuation patterns in zoobenthos populations as a key to underlying factors: Climatic or man-induced. *ICES Journal of Marine Science*, 53, 964-971.
- Beukema JJ, Honkoop PJC, & Dekker R (1998) Recruitment in *Macoma balthica* after mild and cold winters and its possible control by egg production and shrimp predation. *Hydrobiologia*, 375/376, 23-34.
- Beukema JJ, Drent J, & Honkoop PJC (2001) Maximizing lifetime egg production in a Wadden Sea population of the tellinid bivalve *Macoma balthica*: a trade-off between immediate and future reproductive outputs. *Marine Ecology Progress Series*, 209, 119-129.
- Beukema JJ & Meehan BW (1985) Latitudinal variation in linear growth and other shell characteristics of *Macoma balthica*. *Marine Biology*, 90, 27-33.
- Beverton R-JH (1995) Spatial limitation of population size: The concentration hypothesis. *Netherlands Journal of Sea Research*, 34, 1-6.
- Boddeke R (1976) The seasonal migration of the brown shrimp *Crangon crangon*. *Netherlands Journal of Sea Research*, 10, 102-130.
- Boddeke R (1996) Changes in the brown shrimp (*Crangon crangon* L.) population off the Dutch coast in relation to fisheries and phosphate discharge. *ICES Journal of Marine Science*, 53, 995-1002.
- Boehlert GW & Mundy BC (1988) Roles of behavioral and physical factors in larval and juvenile fish recruitment to estuarine nursery areas. *American Fisheries Society Symposium*, 3, 51-67.
- Bond AB & Kamil AC (1998) Apostatic selection by blue jays produces balanced polymorphism in virtual prey. *Nature*, 395, 594-596.
- Bonsdorff E, Norkko A, and Boström C (1993) Recruitment and population maintenance of the bivalve *Macoma balthica* (L.). Factors affecting settling success and early survival on shallow sandy bottoms. *Biology and Ecology of Shallow Coastal Waters*. (eds. A. Eleftheriou, A. D. Ansell, & C. J. Smith), Olsen & Olsen Publ., Fredensborg, Denmark.
- Bonsdorff E, Norkko A, & Sandberg E (1995) Structuring zoobenthos: The importance of predation, siphon cropping and physical disturbance. *Journal of Experimental Marine Biology and Ecology*, 192, 125-144.
- Bouma H, Duiker JMC, De Vries PP, Herman PMJ, & Wolff WJ (2001) Spatial pattern of early recruitment of *Macoma balthica* (L.) and *Cerastoderma edule* (L.) in relation to sediment dynamics on a highly dynamic intertidal sandflat. *Journal of Sea Research*, 45, 79-93.
- Bourget E (1983) Seasonal variations of cold tolerance in intertidal molluscs and relation to environmental conditions in the St. Lawrence estuary. *Canadian Journal of Zoology*, 61, 1193-1201.
- Broekhuysen GJ (1936) On development, growth and distribution of *Carcinides maenas* (L.). *Archives Néerlandais de Zoologie*, 3, 156-164.
- Buchanan S & Babcock R (1997) Primary and secondary settlement by the greenshell mussel *Perna canaliculus*. *Journal of Shellfish Research*, 16, 71-76.
- Bunskoeke EJ, Ens BJ, Hulscher JB, & De Vlas SJ (1996) Why do oystercatchers *Haematopus ostralegus* switch from feeding on Baltic tellin *Macoma balthica* to feeding on the ragworm *Nereis diversicolor* during the breeding season? *Ardea*, 84A, 91-104.
- Burch A & Seed R (2000) Foraging behaviour of *Carcinus maenas* on *Mytilus edulis*: the importance of prey presentation. *Journal of the Marine Biological Association of the United Kingdom*, 80, 799-810.
- Burke JS, Tanaka M, & Seikai T (1995) Influence of light and salinity on behaviour of larval Japanese flounder (*Paralichthys olivaceus*) and implications for inshore migration. *Netherlands Journal of Sea Research*, 34, 59-69.
- Caceres Martinez J, Robledo JAF, & Figueras A (1993) Settlement of mussels *Mytilus galloprovincialis* on an exposed rocky shore in Ria de Vigo, northwestern Spain. *Marine Ecology Progress Series*, 93, 195-198.
- Caceres Martinez J, Robledo JAF, & Figueras A (1994) Settlement and post-larvae behaviour of *Mytilus galloprovincialis*: Field and laboratory experiments. *Marine Ecology Progress Series*, 112, 107-117.
-

- Cadée GC (1976) Sediment reworking by *Arenicola marina* on tidal flats in the Dutch Wadden Sea. *Netherlands Journal of Sea Research*, 10, 331-353.
- Cadée GC & Hegeman J (1977) Distribution of primary production of benthic microflora living on tidal flats in the Dutch Wadden Sea. *Netherlands Journal of Sea Research*, 11, 24-41.
- Commito JA (1982) Importance of predation by infaunal polychaetes in controlling the structure of a soft-bottom community in Maine, USA. *Marine Biology*, 68, 77-81.
- Commito JA & Shrader PB (1985) Benthic community response to experimental additions of the polychaete *Nereis virens*. *Marine Biology*, 86, 101-107.
- Crawley MJ (1993) *GLIM for ecologists*. Blackwell Scientific Publications, Oxford.
- Creutzberg F (1961) The orientation of migrating elvers (*Anguilla anguilla* Turt.) in a tidal area. *Netherlands Journal of Sea Research*, 1, 257-338.
- Creutzberg F, Eltink ATGW, and Van Noort GJ (1978) The migration of plaice larvae *Pleuronectes platessa* into the Western Wadden Sea. *Physiology and behaviour of marine organisms*. (eds. D. S. McLusky & A. J. Berry), pp. 243-251 Pergamon Press, Oxford.
- Cummings VJ, Pridmore RD, Thrush SF, & Hewitt JE (1996) Effect of the spionid polychaete *Boccardia syrtis* on the distribution and survival of juvenile *Macomona liliana* (Bivalvia: Tellinacea). *Marine Biology*, 126, 91-98.
- Curtis LA (1990) Parasitism and the movements of intertidal gastropod individuals. *Biological Bulletin*, 179, 105-112.
- Dahlgren CP & Eggleston DB (2000) Ecological processes underlying ontogenetic habitat shifts in a coral reef fish. *Ecology*, 81, 2227-2240.
- Dankers N & Binsbergen MA (1984) The polychaete *Nereis diversicolor* is also migrating (In Dutch). *De Levende Natuur*, 1984, 14-15.
- De Blok JW & Tan-Maas M (1977) Function of byssus threads in young post-larval *Mytilus*. *Nature*, 267, 558-558.
- De Boer, M. (1978) Waterbeweging in aanlopen en buitendelta's van de zeegaten van de Waddenzee. Notitie 78H248: Rijkswaterstaat, Directie Noord-Holland, studie dienst Hoorn, Hoorn
- De Jong J & Koks B (1999) Wadvogels in de westelijke Noordpolder. *Grauwe gors*, 27, 117-134.
- De Jonge VN & Van den Bergs J (1987) Experiments on the resuspension of estuarine sediments containing benthic diatoms. *Estuarine Coastal and Shelf Science*, 24, 725-740.
- De Montaudouin X (1997) Potential of bivalves' secondary settlement differs with species: A comparison between cockle (*Cerastoderma edule*) and clam (*Ruditapes philippinarum*) juvenile resuspension. *Marine Biology*, 128, 639-648.
- De Robertis A, Jaffe JS, & Ohman MD (2000) Size-dependent visual predation risk and the timing of vertical migration in zooplankton. *Limnology and Oceanography*, 45, 1838-1844.
- De Vlas, J. (1981) Effectenstudie pijpleiding F3-Eemshaven. Voorselectie van 3 nader te onderzoeken trace's door de Waddenzee. Report Rijksinstituut voor Natuurbeheer, Den Burg pp 1-38
- De Wolf P (1973) Ecological observations on the mechanisms of dispersal of barnacle larvae during planktonic life and settling. *Netherlands Journal of Sea Research*, 6, 1-129.
- Dean D (1978) Migration of the sandworm *Nereis virens* during winter nights. *Marine Biology*, 45, 165-173.
- Dekinga A & Piersma T (1993) Reconstructing diet composition on the basis of faeces in a mollusc-eating wader, the knot *Calidris canutus*. *Bird Study*, 40, 144-156.
- Dekker, R. and De Bruin, W. (1998) Het macrozoobenthos op twaalf raaien in de Waddenzee en de Eems-Dollard in 1997. 1998-3: pp 1-53 Netherlands Institute for Sea Research,
- del Norte-Campus AGC & Temming A (1994e) Daily activity, feeding and rations in gobies, and brown shrimp in the northern Wadden Sea. *Marine Ecology Progress Series*, 115, 41-53.
- Desroy N, Retiere C, & Thiebaut E (1998) Infaunal predation regulates benthic recruitment: an experimental study of the influence of the predator *Nephtys hombergii* (Savigny) on recruits of *Nereis diversicolor* (O.F. Müller). *Journal of*

- 
- Experimental Marine Biology and Ecology*, 228, 257-272.
- Devries MC, Tankersley RA, Forward RBJ, Kirby-Smith WW, & Luettich RAJ (1994) Abundance of estuarine crab larvae is associated with tidal hydrologic variables. *Marine Biology*, 118, 403-413.
- Dingle H (1996) *Migration. The biology of life on the move*. Oxford University Press, New York Oxford.
- Duineveld, G. C. A. (1992c) The macrobenthic fauna in the Dutch sector of the North Sea in 1991.1992-6: pp 1-55 Netherlands Institute for Sea Research, Den Burg
- Duineveld, G. C. A. and Belgers, J. J. M. (1993) The macrobenthic fauna in the Dutch sector of the North Sea in 1992. NIOZ-Report 1993-11: pp 1-55 Netherlands Institute for Sea Research, Den Burg
- Durell SEIVd & Kelly CP (1990) Diets of dunlin *Calidris alpina* and grey plover *Pluvialis squatarola* on the Wash as determined by dropping analysis. *Bird Study*, 37, 44-47.
- Edelaar, P. (1998) Evolution of anti-predation traits: dynamic interactions between individuals and populations.3: pp 1-41 CEES, Groningen
- Elmgren R, Ankar S, Marteleur B, & Ejdung G (1986) Adult interference with postlarvae in soft-bottom sediments: the *Pontoporeia-M. balthica* example. *Ecology*, 67, 827-836.
- Emerson CW & Grant J (1991) The control of soft-shell clam (*Mya arenaria*) recruitment on intertidal sandflats by bedload sediment transport. *Limnology and Oceanography*, 36, 1288-1300.
- Essink, K. (1978) The effects of pollution by organic waste on macrofauna in the Eastern Dutch Wadden Sea. NIOZ-Report 1: pp 1-135 Netherlands Institute for Sea Research, Den Burg
- Essink K, Eppinga J, & Dekker R (1998a) Long-term changes (1977-1994) in intertidal macrozoobenthos of the Dollard (Ems Estuary) and effects of introduction of the North American spionid polychaete *Marenzelleria cf. wireni*. *Senckenbergiana Maritima*, 28, 221-225.
- Essink K, Beukema JJ, Madsen PB, Michaelis H, & Vedel GR (1998b) Long-term development of biomass of intertidal macrozoobenthos in different parts of the Wadden Sea. Governed by nutrient loads? *Senckenbergiana Maritima*, 29, 25-35.
- Essink K & Kleef HL (1993) Distribution and life cycle of the North American spionid polychaete *Marenzelleria viridis* (Verrill, 1873) in the Ems estuary. *Netherlands Journal of Aquatic Ecology*, 27, 237-246.
- Essink K, Kleef HL, & Visser W (1989) On the pelagic occurrence and dispersal of the benthic amphipod *Corophium volutator*. *Journal of the Marine Biological Association of the United Kingdom*, 69, 11-15.
- Farke H & Berghuis EM (1979) Spawning, larval development and migration behaviour of *Arenicola marina* in the laboratory. *Netherlands Journal of Sea Research*, 13, 512-528.
- Farke H, De Wilde PAWJ, & Berghuis EM (1979) Distribution of juvenile and adult *Arenicola marina* on a tidal mud flat and the importance of nearshore areas for recruitment. *Netherlands Journal of Sea Research*, 13, 354-361.
- Fernandez TF, Huxham M, & Piper SR (1999) Predator caging experiments: a test of the importance of scale. *Journal of Experimental Marine Biology and Ecology*, 241, 137-154.
- Flach E and Tamaki A (2001) Competitive bioturbators on intertidal sand flats in the European Wadden Sea and Ariake Sound in Japan. *Ecological comparisons of sedimentary shores*. (ed. K. Reise), pp. 149-168 Springer, Berlin.
- Flach EC (1992a) Disturbance of benthic infauna by sediment-reworking activities of the lugworm *Arenicola marina*. *Netherlands Journal of Sea Research*, 30, 81-89.
- Flach EC (1992b) The influence of four macrozoobenthic species on the abundance of the amphipod *Corophium volutator* on tidal flats. *Netherlands Journal of Sea Research*, 29, 379-394.
- Flach EC (1993) The distribution of the amphipod *Corophium arenarium* in the Dutch Wadden Sea: Relationships with sediment composition and the presence of cockles and lugworms. *Netherlands Journal of Sea Research*, 31, 281-290.
- Flach EC (1996) The influence of the cockle, *Cerastoderma edule*, on the macrozoobenthic community of tidal flats in the Wadden Sea. *Marine Ecology*, 17, 87-98.
-

- Flach EC & Beukema JJ (1994) Density-governing mechanisms in populations of the lugworm *Arenicola marina* on tidal flats. *Marine Ecology Progress Series*, 115, 139-149.
- Flach EC & De Bruin W (1993) Effects of *Arenicola marina* and *Cerastoderma edule* on distribution, abundance and population structure of *Corophium volutator* in Gullmarsfjorden western Sweden. *Sarsia*, 78, 105-118.
- Flach EC & De Bruin W (1994) Does the activity of cockles, *Cerastoderma edule* (L.) and lugworms, *Arenicola marina* L., make *Corophium volutator* Pallas more vulnerable to epibenthic predators: A case of interaction modification? *Journal of Experimental Marine Biology and Ecology*, 182, 265-285.
- Fonds M (1973) Sand gobies (Pomatoschistus, Gobiidae) of the Dutch Wadden Sea. *Netherlands Journal of Sea Research*,
- Ford RB & Paterson DM (2001) Behaviour of *Corophium volutator* in still versus flowing water. *Estuarine Coastal and Shelf Science*, 52, 357-362.
- Forward RBJ & Tankersley RA (2001) Selective tidal-stream transport of marine animals. *Oceanography and Marine Biology: an annual review*, 39, 305-353.
- Forward RB, Jr. & Rittschof D (2000) Alteration of photoresponses involved in diel vertical migration of a crab larva by fish mucus and degradation products of mucopolysaccharides. *Journal of Experimental Marine Biology and Ecology*, 245, 277-292.
- Galaktionov KV (1992) Seasonal dynamics of the age composition of hemipopulations of daughter sporocysts of pygmaeus microphallids (Trematoda: Microphallidae) in intertidal molluscs *Littorina saxatilis* from the Barents Sea. *Parazitologiya*, 26, 462-469.
- Gibson RN (1994) Impact of habitat quality and quantity on the recruitment of juvenile flatfishes. *Netherlands Journal of Sea Research*, 32, 191-206.
- Goethe F (1961) The moult gatherings and moult migration of the Shelduck in north-west Germany. *British Birds*, 54, 145-161.
- Grant J (1981) Sediment transport and disturbance on an intertidal sandflat: Infaunal distribution and recolonization. *Marine Ecology Progress Series*, 6, 249-255.
- Gray DR & Hodgson AN (1999) Endogenous rhythms of locomotor activity in the high-shore limpet, *Helcion pectunculus* (Patellogastropoda). *Animal Behaviour*, 57, 387-391.
- Gray JS (1939) Studies in animal locomotion. VIII. The kinetics of locomotion of *Nereis diversicolor*. *Journal of Experimental Biology*, 16, 9-18.
- Green RH (1973) Growth and mortality in an arctic population of *Macoma balthica* (Pelecypoda, Tellinidae). *Journal of the Fisheries Research Board of Canada*, 30, 1345-1348.
- Greenwood-Lee JM & Taylor PD (2001) The evolution of dispersal in spatially varying environments. *Evolutionary Ecology Research*, 3, 649-665.
- Gregory RS & Levings CD (1996) The effects of turbidity and vegetation on the risk of juvenile salmonids, *Oncorhynchus* spp., to predation by adult cutthroat trout, *O. clarkii*. *Environmental Biology of Fishes*, 47, 279-288.
- Gregory RS & Levings CD (1998) Turbidity reduces predation on migrating juvenile Pacific salmon. *Transactions of the American Fisheries Society*, 127, 275-285.
- Guillou J & Tartu C (1994) Post-larval and juvenile mortality in a population of the edible cockle *Cerastoderma edule* (L.) from northern Brittany. *Netherlands Journal of Sea Research*, 33, 103-111.
- Günther CP (1991) Settlement of *Macoma balthica* on an intertidal sandflat in the Wadden Sea. *Marine Ecology Progress Series*, 76, 73-79.
- Günther CP (1992) Dispersal of intertidal invertebrates: a strategy to react to disturbances of different scales? *Netherlands Journal of Sea Research*, 30, 45-56.
- Günther CP and Niesel V (1999) Effects of the ice winter 1995/96. *The Wadden Sea ecosystem. Stability properties and mechanisms*. (ed. S. Dittman), pp. 193-205 Springer-Verlag, Berlin.
- Hanski I (1999) *Metapopulation ecology*. Oxford university press, Oxford.
-

- 
- Hamilton WD (1980) Sex versus non-sex versus parasite. *Oikos*, 35, 282-290.
- Hamilton WD & May RM (1977) Dispersal in a stable habitat. *Nature*, 269, 578-581.
- Hartman-Schröder G (1971) *Annelida, Borstenwürmer, Polychaeta*. Gustav Fisher Verlag, Jena.
- Harvey M & Vincent B (1990) Density, size distribution, energy allocation and seasonal variations in shell and soft tissue growth at two tidal levels of a *Macoma balthica* (L.) population. *Journal of Experimental Marine Biology and Ecology*, 142, 151-168.
- Harvey M & Vincent B (1991) Spatial variability of length-specific production in shell, somatic tissue and sexual products of *Macoma balthica* in the Lower St. Lawrence Estuary: I. Small and meso scale variability. *Marine Ecology Progress Series*, 75, 55-66.
- Harvey M, Vincent B, & Gratton Y (1993) Spatial variability of length-specific production in shell, somatic tissue and sexual products of *Macoma balthica* in the Lower St. Lawrence Estuary: II. Large scale variability. *Marine Biology*, 115, 421-433.
- Hecht T & Van Der Lingen CD (1992) Turbidity-induced changes in feeding strategies of fish in estuaries. *South African Journal of Zoology*, 27, 95-107.
- Holling CS (1965) The functional response of predators to prey density and its role in mimicry and population regulation. *Memoirs of the Entomological Society of Canada*, 45, 1-60.
- Holtmann, S. E., Duineveld, G. C. A., Mulder, M., and De Wilde, P. A. W. J. (1995) The macrobenthic fauna in the Dutch sector of the North Sea in 1994 and a comparison with previous data. NIOZ-Report 1995-7: pp 1-103 Netherlands Institute for Sea Research, Den Burg
- Holtmann, S. E., Duineveld, G. C. A., Mulder, M., and De Wilde, P. A. W. J. (1996a) The macrobenthic fauna in the Dutch sector of the North Sea in 1995 and a comparison with previous data. NIOZ-Report 1996-5: pp 1-103 Netherlands Institute for Sea Research, Den Burg
- Holtmann SE, Groenewold A, Schrader KHM, Asjes J, Craeymeersch JAM, Duineveld GCA, Van Bostelen AJ & Van der Meer J (1996b) *Atlas of the zoobenthos of the Dutch continental shelf*. Ministry of Transport, Public Works and Water Management, North Sea Directorate, Rijswijk.
- Holtmann, S. E., Duineveld, G. C. A., Mulder, M., and De Wilde, P. A. W. J. (1997) The macrobenthic fauna in the Dutch sector of the North Sea in 1996 and a comparison with previous data. NIOZ-Report 1997-8: pp 1-103 Netherlands Institute for Sea Research, Den Burg
- Holtmann, S. E., Duineveld, G. C. A., Mulder, M., and De Wilde, P. A. W. J. (1998) The macrobenthic fauna in the Dutch sector of the North Sea in 1997 and a comparison with previous data. NIOZ-Report 1998-5: pp 1-103 Netherlands Institute for Sea Research, Den Burg
- Holtmann, S. E., Duineveld, G. C. A., and Mulder, M. (1999) The macrobenthic fauna in the Dutch sector of the North Sea in 1998 and a comparison with previous data. NIOZ-Report 1999-5: pp 1-105 Netherlands Institute for Sea Research, Den Burg
- Honkoop PJC & Van der Meer J (1997) Reproductive output of *Macoma balthica* populations in relation to winter-temperature and intertidal-height mediated changes of body mass. *Marine Ecology Progress Series*, 149, 155-162.
- Honkoop PJC, Van der Meer J, Beukema JJ, & Kwast D (1998) Does temperature-influenced egg production predict the recruitment in the bivalve *Macoma balthica*? *Marine Ecology Progress Series*, 164, 229-235.
- Hostens K (2000) Spatial patterns and seasonality in the epibenthic communities of the Westerschelde (Southern Bight of the North Sea). *Journal of the Marine Biological Association of the United Kingdom*, 80, 27-36.
- Hostens K & Mees J (1999) The mysid-feeding guild of demersal fishes in the brackish zone of the Westerschelde estuary. *Journal of Fish Biology*, 55, 704-719.
- Hughes DA (1969) Response to salinity change as a tidal transport mechanism of pink shrimp *Penaeus duorarum*. *Behaviour*, 136, 43-53.
- Hughes RG & Gerdol V (1997) Factors affecting the distribution of the amphipod *Corophium volutator* in two estuaries in south-east England. *Estuarine Coastal and Shelf Science*, 44, 612-627.
- Hughes RG & Horsfall IM (1990) Differences in the swimming behavior of the amphipod *Corophium volutator* from different populations. *Journal of the Marine Biological Association of the United Kingdom*, 70, 143-148.
-



- Hulscher JB (1973) Burying-depth and trematode infection in *Macoma balthica*. *Netherlands Journal of Sea Research*, 6, 141-156.
- Hulscher JB (1982) The oystercatcher *Haematopus ostralegus* as a predator of the bivalve *Macoma balthica* in the Dutch Wadden Sea. *Ardea*, 70, 89-152.
- Hummel H, Bijok P, & Bogaards RH (1996) Effects of tidal zonation on size and genetic traits of *Mytilus edulis* (L.) and *Macoma balthica* (L.). *Polskie Archiwum Hydrobiologii*, 43, 431-445.
- Hunt HL & Mullineaux LS (2002) The roles of predation and postlarval transport in recruitment of the soft shell clam (*Mya arenaria*). *Limnology and Oceanography*, 47, 151-164.
- Hunt HL & Scheibling RE (1997) Role of early post-settlement mortality in recruitment of benthic marine invertebrates. *Marine Ecology Progress Series*, 155, 269-301.
- Ibing J & Theede H (1975) Zur Gefrierresistenz litoraler Mollusken von der deutschen Nordseeküste. *Kieler Meeresforschung*, 31, 44-49.
- Iles TC & Beverton R-JH (2000) The concentration hypothesis: The statistical evidence. *ICES Journal of Marine Science*, 57, 216-227.
- Ivlev VS (1961) *Experimental biology of feeding in fishes*. Yale University Press, New Haven.
- Jager Z (1999) Selective tidal stream transport of flounder larvae (*Platichthys flesus* L.) in the Dollard (Ems estuary). *Estuarine Coastal and Shelf Science*, 49, 347-362.
- Jaklin S & Günther CP (1996) Macrobenthic driftfauna of the Gröninger plate. *Senckenbergiana Maritima*, 26, 127-134.
- Jaquet N & Raffaelli D (1989) The ecological importance of the sand goby *Pomatoschistus minutus* (Pallas). *Journal of Experimental Marine Biology and Ecology*, 128, 147-156.
- Jensen KT (1992) Dynamics and growth of the cockle, *Cerastoderma edule*, on an intertidal mud-flat in the Danish Wadden Sea: Effects of submersion time and density. *Netherlands Journal of Sea Research*, 28, 335-345.
- Kafanov AI, Danilin DD, & Moschenko AV (1999) Intrapopulation variability of meristic characters in discrete color morphs of the bivalve *Macoma balthica*. *Biologiya Morya*, 25, 13-18.
- Kaiser MJ, Hughes RN, & Reid DG (1990) Chelal morphometry, prey-size selection and aggressive competition in green and red forms of *Carcinus maenas* (L.). *Journal of Experimental Marine Biology and Ecology*, 140, 121-134.
- Kennedy AD (1993) Minimal predation upon meiofauna by endobenthic macrofauna in the Exe Estuary, south west England. *Marine Biology*, 117, 311-319.
- Kermack DM (1955) The anatomy and physiology of the gut of the polychaete *Arenicola marina*. *Proceedings of the Zoological Society of London*, 125, 347-381.
- Keus B (1986) De predatie van de Garnaal (*Crangon crangon*) op het broed van het Nonnetje (*Macoma balthica*). *Interne Verslagen Nederlands Instituut voor Onderzoek der Zee*, 1986, 1-43.
- Kim WS, Huh HT, Huh SH, & Lee TW (2001) Effects of salinity on endogenous rhythm of the Manila clam, *Ruditapes philippinarum* (Bivalvia: Veneridae). *Marine Biology*, 138, 157-162.
- King M (1995) *Fisheries biology, assessment and management*. Fishing news books, Oxford.
- Klein Breteler WCM (1976) Settlement, growth and production of the shore crab *Carcinus maenas* on tidal flats of the Dutch Wadden Sea. *Netherlands Journal of Sea Research*, 10, 354-376.
- Korringa P (1947) Relations between the moon and periodicity in the breeding of marine animals. *Ecological Monographs*, 17, 347-381.
- Korringa P (1950) Wie let in april eens op de grote groene zeeduizendpoot? *Natura*, 47, 53
- Kuipers BR & Dapper R (1984) Nursery function of Wadden Sea tidal flats for the brown shrimp *Crangon crangon*. *Marine Ecology Progress Series*, 17, 171-181.
- Kullberg C, Fransson T, & Jakobsson S (1996) Impaired predator evasion in fat blackcaps (*Sylvia atricapilla*). *Proceedings of the Royal Society of London Series B Biological Sciences*, 263, 1671-1675.
-

- 
- Kühl H (1970) Nahrungsuntersuchungen am Stint (*Osmerus eperlanus* L.) im Elbe-Mündungsgebiet. *Archiv für Fischerei Wissenschaft*, 21, 222-231.
- Lane DJW, Beaumont AR, & Hunter JR (1985) Byssus drifting and the drifting threads of the young post-larval mussel *Mytilus edulis*. *Marine Biology*, 84, 301-308.
- Lane DJW, Nott JA, & Crisp JD (1982) Enlarged stem glands in the foot of the post-larval mussel *Mytilus edulis*: adaptation for bysso-pelagic migration. *Journal of the Marine Biological Association of the United Kingdom*, 62, 809-818.
- Lauckner G (1983) Diseases of Mollusca: Bivalvia. *Diseases of Marine Animals. Volume II: Introduction, Bivalvia to Scaphopoda*. pp. 632-761 Biologische Anstalt Helgoland, Hamburg.
- Lauckner G (1986) Ecological effects of larval trematode infestation on littoral marine invertebrate populations. *Proceedings of the International congress of Parasitology*. pp. 391-398 Brisbane.
- Lawrie SM & Raffaelli DG (1998) In situ swimming behaviour of the amphipod *Corophium volutator* (Pallas). *Journal of Experimental Marine Biology and Ecology*, 224, 237-251.
- Lim SSL & Green RH (1991) The relationship between parasite load, crawling behaviour, and growth rate of *Macoma balthica* (L.) (Mollusca, Pelecypoda) from Hudson Bay, Canada. *Canadian Journal of Zoology*, 69, 2202-2208.
- Lindeboom, H. J., Bergman, M. J. N., and De Gee, A. (1996) Ecologisch minst kwetsbare locaties voor exploratieboringen in de Waddenzee en Noordzee kustzone. NIOZ-Report 1996-4: pp 1-116 Netherlands Institute for Sea Research, Den Burg
- Little C & Nix W (1976a) The burrowing and floating behaviour of the gastropod *Hydrobia ulvae*. *Estuarine and Coastal Marine Science*, 4, 537-544.
- Lorenzen CJ (1967) Determination of chlorophyll and pheo-pigments: spectrophotometric equations. *Limnology and Oceanography*, 12, 343-346.
- Louters, T. and Gerritsen, F. (1994) The riddle of the sands. A tidal system's answer to a rising sea level. RIKZ-report 94.040: pp 1-69, Den Haag
- Martel A & Chia FS (1991) Foot-raising behavior and active participation during the initial phase of post-metamorphic drifting in the gastropod *Lacuna* spp. *Marine Ecology Progress Series*, 72, 247-254.
- Mascaro M & Seed R (2001) Choice of prey size and species in *Carcinus maenas* (L.) feeding on four bivalves of contrasting shell morphology. *Hydrobiologia*, 449, 159-170.
- Masson S, Desrosiers G, & Retiere C (1995) Périodicité d'alimentation du polychète *N. diversicolor* (O.F. Müller) selon les changements de la marée. *Ecoscience*, 2, 20-27.
- Matthiessen GC (1960) Intertidal zonation in population of *Mya arenaria*. *Limnology and Oceanography*, 5, 381-388.
- McIntosh AR & Townsend CR (1995) Contrasting predation risks presented by introduced brown trout and native common river galaxias in New Zealand streams. *Canadian Journal Of Fisheries And Aquatic Sciences*, 52, 1821-1833.
- McPeck MA & Holt RD (1992) The evolution of dispersal in spatially and temporally varying environments. *American Naturalist*, 140, 1010-1027.
- Miron GY & Desrosiers GL (1990) Distributions and population structures of two intertidal estuarine polychaetes in the lower St. Lawrence estuary (Quebec, Canada) with special reference to environmental factors. *Marine Biology*, 105, 297-306.
- Moment GB (1962) Reflexive selection: a possible answer to an old puzzle. *Science*, 136, 262-263.
- Moreira F (1994a) Diet and feeding rates of knots *Calidris canutus* in the Tagus estuary (Portugal). *Ardea*, 82, 133-135.
- Moreira F (1994b) Diet, prey-sized selection and intake rates of Black-tailed Godwits *Limosa limosa* feeding on mudflats. *Ibis*, 136, 349-355.
- Mouritsen KN (1997) Crawling behaviour in the bivalve *Macoma balthica*: The parasite-manipulation hypothesis revisited. *Oikos*, 79, 513-520.
- Nash RDM, Santos RS, Geffen AJ, Hughes G, & Ellis TR (1994) Diel variability in catch rate of juvenile flatfish on two small nursery grounds (Port Erin Bay, Isle of Man and Porto Pim Bay, Faial, Azores). *Journal of Fish Biology*, 44, 35-45.
-

- Newell R (1962) Behavioural aspects of the ecology of *Peringia* (= *Hydrobia*) *ulvae* (Pennant) (Gasteropoda, Prosobranchia). *Proceedings of the Zoological Society of London*, 138, 49-75.
- Niesel V (1999) Study area: the backbarrier tidal flats of Spiekeroog. Meteorological conditions in the East Frisian Wadden Sea from 1994-1996. *The Wadden Sea ecosystem. Stability properties and mechanisms*. (ed. S. Dittman), pp. 19-25 Springer-Verlag, Berlin.
- Nilsson DE (1994) Eyes as optical alarm systems in fan worms and ark clams. *Philosophical Transactions of the Royal Society of London B Biological Sciences*, 346, 195-212.
- Olafsson E, Peterson CH, & Ambrose WG (1994) Does recruitment limitation structure populations and communities of macro-invertebrates in marine soft sediments: the relative significance of pre- and post settlement processes. *Oceanography and Marine Biology: an annual review*, 32, 65-109.
- Olney PJS (1965) Food and feeding habits of the shelduck *Tadorna tadorna*. *Ibis*, 107, 527-532.
- Owen DF & Whiteley D (1986) Reflexive selection: Moment's hypothesis resurrected. *Oikos*, 47, 117-120.
- Owen DF & Whiteley D (1988) The beach clams of Thessaloniki: reflexive or apostatic selection. *Oikos*, 51, 253-255.
- Pekkarinen M (1987) Notes on a gymnophallid trematode assumed to be *Parvatrema affinis* (Jameson and Nicoll, 1913), from *Macoma balthica*. *Annales Zoologici Fennici*, 24, 29-37.
- Piersma T, Hoekstra R, Dekinga A, Koolhaas A, Wolf P, Battley P, & Wiersma P (1993) Scale and intensity of intertidal habitat use by knots *Calidris canutus* in the western Wadden Sea in relation to food, friends and foes. *Netherlands Journal of Sea Research*, 31, 331-357.
- Piersma T, Verkuil Y, & Tulp I (1994) Resources for long-distance migration of knots *Calidris canutus islandica* and *C. c. canutus*: How broad is the temporal exploitation window of benthic prey in the western and eastern Wadden Sea? *Oikos*, 71, 393-407.
- Pihl L & Rosenberg R (1984) Food selection and consumption of the shrimp *Crangon crangon* in some shallow marine areas in western Sweden. *Marine Ecology Progress Series*, 15, 159-168.
- Plante CJ & Mayer LM (1994) Distribution and efficiency of bacteriolysis in the gut of *Arenicola marina* and three additional deposit feeders. *Marine Ecology Progress Series*, 109, 183-194.
- Plante CJ & Mayer LM (1996) Seasonal variability in the bacteriolytic capacity of the deposit feeder *Arenicola marina*: Environmental correlates. *Aquatic Microbial Ecology*, 11, 101-109.
- Poutsma, J. (1969) Oriënterend onderzoek naar het menu van de Bergeend (*Tadorna tadorna* (Linnaeus)) en de Bonte strandloper (*Calidris alpina* (Linnaeus)) op het Groninger Wad. Report Zoologisch laboratorium, Baffo pp 1-38.
- Pulfrich A (1996) Attachment and settlement of post-larval mussels (*Mytilus edulis* L.) in the Schleswig-Holstein, Wadden Sea. *Journal of Sea Research*, 36, 239-250.
- Ratcliffe PJ, Jones NV, and Walters NJ (1981) The survival of *Macoma balthica* (L.) in mobile sediments. *Feeding and survival strategies of estuarine organisms*. (eds. N. V. Jones & W. J. Wolff), pp. 91-108 Plenum Press, New York and London.
- Reise K (1978) Experiments on epibenthic predation in the Wadden Sea. *Helgoländer Meeresuntersuchungen*, 31, 55-101.
- Reise K (1979a) Moderate predation on meiofauna by the macrobenthos of the Wadden Sea. *Helgoländer Meeresuntersuchungen*, 32, 453-465.
- Reise K (1979b) Spatial configurations generated by motile benthic polychaetes. *Helgoländer Meeresuntersuchungen*, 32, 55-72.
- Reise K (1985) *Tidal flat ecology*. Springer-Verlag, Berlin.
- Retraubun ASW, Dawson M, & Evans SM (1996) The role of the burrow funnel in feeding processes in the lugworm *Arenicola marina* (L.). *Journal of Experimental Marine Biology and Ecology*, 202, 107-118.
- Richards M, Edwards F, & Huxham M (2002) The effects of the adult density of *Macoma balthica* on the recruitment of juvenile bivalves: a field experiment. *Journal of Sea Research*, 47, 54-54.
- Richards MG, Huxham M, & Bryant A (1999) Predation: a causal mechanism for variability in intertidal bivalve populations.
-

- 
- Journal of Experimental Marine Biology and Ecology*, 241, 159-177.
- Ridderinkhof H, Zimmerman JTF, & Philippart ME (1990) Tidal exchange between the North Sea and Dutch Wadden Sea and mixing time scales of tidal basins. *Netherlands Journal of Sea Research*, 25, 331-350.
- Riisgard HU, Vedel A, & Boye H (1992) Filter-net structure and pumping activity in the polychaete *Nereis diversicolor*. Effects of temperature and pump-modelling. *Marine Ecology Progress Series*, 83, 79-89.
- Ronn C, Bonsdorff E, & Nelson W (1988) Predation as a mechanism of interference within infauna in shallow brackish water soft bottoms, experiments with an infaunal predator *Nereis diversicolor*. *Journal of Experimental Marine Biology and Ecology*, 116, 143-157.
- Rothlisberg PC, Church JA, & Fandry CB (1995) A mechanism for near-shore concentration and estuarine recruitment of post-larval *Penaeus plebejus* Hess (Decapoda, Penaeidae). *Estuarine Coastal and Shelf Science*, 40, 115-138.
- Sale PF (1969) A suggested mechanism for habitat selection by the juvenile manini, *Acanthurus triostegus sandvicensis*. *Behaviour*, 35, 27-44.
- Sanchez-Salazar ME, Griffiths CL, & Seed R (1987a) The effect of size and temperature on the predation of cockles *Cerastoderma edule* (L.) by the shore crab *Carcinus maenas* (L.). *Journal of Experimental Marine Biology and Ecology*, 111, 181-193.
- Sanchez-Salazar ME, Griffiths CL, & Seed R (1987b) The interactive roles of predation and tidal elevation in structuring populations of the edible cockle, *Cerastoderma edule*. *Estuarine Coastal and Shelf Science*, 25, 245-260.
- Sarvala J (1971) Ecology of *Harmothoe sarsi* (Malmgren) (Polychaeta, Polynoidae) in the northern Baltic area. *Annales Zoologici Fennici*, 8, 231-309.
- Scherer B & Reise K (1981) Significant predation on micro- and macrobenthos by the crab *Carcinus maenas*. *Kieler Meeresforschung*, 5, 490-500.
- Senner SE, Norton DW, & West GC (1989) Feeding ecology of western sandpipers, *Calidris mauri*, and dunlins, *Calidris alpina*, during spring migration at Harney Bay, Alaska (USA). *Canadian Field-Naturalist*, 103, 372-379.
- Sewell MA (1996) Detection of the impact of predation by migratory shorebirds: An experimental test in the Fraser River estuary, British Columbia (Canada). *Marine Ecology Progress Series*, 144, 23-40.
- Shaffer PL (1983) Population ecology of *Heteromastus filiformis* (Polychaeta: Capitellidae). *Netherlands Journal of Sea Research*, 17, 106-125.
- Sigurdsson JB, Titman CW, & Davis PA (1976) The dispersal of young post-larval bivalve molluscs by byssus threads. *Nature*, 262, 386-387.
- Smith DAS (1975) Polymorphism and selective predation in *Donax faba* Gmeling (Bivalvia, Tellinacea). *Journal of Experimental Marine Biology and Ecology*, 17, 205-219.
- Sokal RR & Rohlf RJ (1990) *Biometry*. Freeman, New York.
- Sola JC (1996) Population dynamics, reproduction, growth, and secondary production of the mud-snail *Hydrobia ulvae* (Pennant). *Journal of Experimental Marine Biology and Ecology*, 205, 49-62.
- Sörlin T (1988) Floating behaviour in the tellinid bivalve *Macoma balthica* (L.). *Oecologia*, 77, 273-277.
- Spaargaren DH (1980) The significance of seawater viscosity for the tidal transport of common shrimps, *Crangon crangon* (L.). *Ophelia*, 19, 145-153.
- Stam A (1981) The food of pelagic fish in the Dollart. *Kieler Meeresforschung*, 5, 282
- Stamhuis EJ, Videler JJ, & De Wilde PAWJ (1998) Optimal foraging in the thalassinidean shrimp *Callinassa subterranea* Improving food quality by grain size selection. *Journal of Experimental Marine Biology and Ecology*, 228, 197-208.
- Stamm M (1995) Zur Ernährung von *Retusa obtusa* (Montagu, 1803) im Wattgebiet des Jadebusens (Gastropoda: Opisthobranchia). *Senckenbergiana Maritima*, 25, 137-145.
- Strasser M & Günther CP (2001) Larval supply of predator and prey: temporal mismatch between crabs and bivalves after a severe winter in the Wadden Sea. *Journal of Sea Research*, 46, 57-67.
-

- Strasser M, Hertlein A, & Reise K (2001) Differential recruitment of bivalve species in the northern Wadden Sea after the severe winter of 1995/96 and of subsequent milder winters. *Helgoland Marine Research*, 55, 182-189.
- Strasser M, Walensky M, & Reise K (1999) Juvenile-adult distribution of the bivalve *Mya arenaria* on intertidal flats in the Wadden Sea: why are there so few year classes? *Helgoland Marine Research*, 53, 45-55.
- Swennen C (1969) Crawling tracks of trematode infected *Macoma balthica* (L.). *Netherlands Journal of Sea Research*, 4, 376-379.
- Swennen C & Ching HL (1974) Observation on the trematode *Parvatrema affinis*, causative agent of crawling tracks of *Macoma balthica*. *Netherlands Journal of Sea Research*, 8, 108-115.
- Thiebaut E, Dauvin JC, & Wang Z (1996) Tidal transport of *Pectinaria koreni* postlarvae (Annelida: Polychaeta) in the Bay of Seine (eastern English Channel). *Marine Ecology Progress Series*, 138, 63-70.
- Thiel M & Reise K (1993) Interaction of nemertines and their prey on tidal flats. *Netherlands Journal of Sea Research*, 31, 163-172.
- Tita G, Desrosiers G, Vincx M, & Nozais C (2000) Predation and sediment disturbance effects of the intertidal polychaete *Nereis virens* (Sars) on associated meiofaunal assemblages. *Journal of Experimental Marine Biology and Ecology*, 243, 261-282.
- Tydemann, P. (2000) Het macrozoobenthos op vijf permanente quadraten op het Groninger Wad in 1999. Report RIKZ/ 2000.039: pp 1-37
- Van de Kam J, Ens B, Piersma T & Zwarts L (1999) *Ecologische atlas van de Nederlandse Wadvogels*. Schuyt & Co, Haarlem.
- Van der Meer J, Piersma T, and Beukema JJ (2001a) Population dynamics of benthic species on tidal flats: the possible roles of shorebird predation. *Ecological comparison of sedimentary shores*. (ed. K. Reise), pp. 317-332 Springer, Berlin.
- Van der Meer J, Beukema JJ, & Dekker R (2001b) Long-term variability in secondary production of an intertidal bivalve population is primarily a matter of recruitment variability. *Journal of Animal Ecology*, 70, 159-169.
- Van der Veer HW, Feller RJ, Weber A, & Witte JIJ (1998) Importance of predation by crustaceans upon bivalve spat in the intertidal zone of the Dutch Wadden Sea as revealed by immunological assays of gut contents. *Journal of Experimental Marine Biology and Ecology*, 231, 139-157.
- Van Scheppingen, Y. and Groenewold, A. (1990) De ruimtelijke verspreiding van het benthos in de zuidelijke Noordzee. De Nederlandse kustzone overzicht 1988-1989. Report 90-003: pp 1-27 Rijkswaterstaat, Directie Noordzee, Dienst getijdewateren., Den Haag
- Vedel A, Andersen BB, & Riisgard HU (1994) Field investigations of pumping activity of the facultatively filter-feeding polychaete *Nereis diversicolor* using an improved infrared phototransducer system. *Marine Ecology Progress Series*, 103, 91-101.
- Venema SC & Creutzberg F (1973) Seasonal migration of the swimming crab *Macropipus holsatus* in an estuarine area controlled by tidal streams. *Netherlands Journal of Sea Research*, 7, 94-102.
- Verrill, A. E. (1873) Report upon the invertebrate animals of Vineyard Sound and adjacent waters, with an account of the physical features of the region. 1871-1872: pp 295-852
- Vincent B, Joly D, & Harvey M (1994) Spatial variation in growth of the bivalve *Macoma balthica* (L.) on a tidal flat: Effects of environmental factors and intraspecific competition. *Journal of Experimental Marine Biology and Ecology*, 181, 223-238.
- Wanink JH & Zwarts L (1993) Environmental effects on the growth rate of intertidal invertebrates and some implications for foraging waders. *Netherlands Journal of Sea Research*, 31, 407-418.
- Warman CG, O' Hare TJ, & Naylor E (1991) Vertical swimming in wave-induced currents as a control mechanism of intertidal migration by a sand-beach isopod. *Marine Biology*, 111, 49-54.
- Warwick RM & Price R (1975) Macrofauna production of an estuarine mud flat. *Journal of the Marine Biological Association of the United Kingdom*, 55, 1-18.
- Wennhage H & Pihl L (2001) Settlement patterns of newly settled plaice (*Pleuronectes platessa*) in a non-tidal Swedish fjord in relation to larval supply and benthic predators. *Marine Biology*, 139, 877-889.

- 
- Werner B (1956) Über die Winterwanderung von *Arenicola marina* L. (Polychaeta sedentaria). *Helgoländer Meeresuntersuchungen*, 5, 353-378.
- Werner EE (1986) Amphibian metamorphosis: growth rate, predation risk, and the optimal size at transformation. *American Naturalist*, 128, 319-341.
- Werner EE & Gilliam JF (1984) The ontogenetic niche and species interactions in size-structured populations. *Annual Review of Ecology and Systematics*, 15, 393-425.
- Wolff WJ & De Wolf L (1977) Biomass and production of zoobenthos in the Grevelingen Estuary, The Netherlands. *Estuarine and Coastal Marine Science*, 5, 1-24.
- Wolff WJ, Van Haperen AMM, Sandee AJJ, Baptist HJM, & Saeijs HLF (1975) The trophic role of birds in the Grevelingen estuary, The Netherlands, as compared to their role in the saline lake Grevelingen. *10th European Symposium on Marine Biology, Ostend, Belgium*, 2, 673-689.
- Yates MG & Goss-Custard JD (1991) A comparison between high water and low water counts of shorebirds on the Wash, east England. *Bird Study*, 38, 179-187.
- Zühlke R & Reise K (1994) Response of macrofauna to drifting tidal sediments. *Helgoländer Meeresuntersuchungen*, 48, 277-289.
- Zwarts L (1991) Seasonal variation in body weight of the bivalves *Macoma balthica*, *Scrobicularia plana*, *Mya arenaria* and *Cerastoderma edule* in the Dutch Wadden Sea. *Netherlands Journal of Sea Research*, 28, 231-245.
- Zwarts L & Blomert AM (1992) Why knot *Calidris canutus* take medium-sized *Macoma balthica* when six prey species are available. *Marine Ecology Progress Series*, 83, 113-128.
- Zwarts L, Blomert AM, & Wanink JH (1992) Annual and seasonal variation in the food supply harvestable by knot *Calidris canutus* staging in the Wadden Sea in late summer. *Marine Ecology Progress Series*, 83, 129-139.
- Zwarts L and Wanink JH (1984) How oystercatchers and curlews successively deplete clams. *Coastal waders and wildfowl in winter*. (eds. J. D. Goss-Custard & W. G. Hale), pp. 69-83 Cambridge University Press, Cambridge.
-

